

Universal Pion Freeze-out Phase-Space Density

D. Ferenc, B. Tomášik, U. Heinz, Inst. f. Theor. Physik, Universität Regensburg

G. Bertsch has indicated [1] a possibility to measure the pion freeze-out phase-space density and thereby test the local thermal equilibrium in a pion source. In case of thermal equilibrium at temperature T , identical pions of energy E would follow the Bose-Einstein distribution

$$f = \frac{1}{e^{\frac{E}{T}} - 1}. \quad (1)$$

An average of this function over different phase-space regions is the quantity to be measured. When the p_T -spectrum is parameterized by an exponential with $T_{\text{eff}}(y)$ being the inverse slope parameter, averaging over the spatial coordinates yields

$$\langle f \rangle(p_T, y) = \frac{\frac{\sqrt{\pi}}{2} \frac{\sqrt{\lambda_{\text{dir}}(p_T, y)}}{E_p T_{\text{eff}}^2(y)} \exp\left(-\frac{p_T}{T_{\text{eff}}(y)}\right) \frac{dn^-}{dy}(y)}{R_s(p_T, y) \sqrt{R_o^2(p_T, y) R_l^2(p_T, y) - R_{ol}^4(p_T, y)}}. \quad (2)$$

This equation comprises information from essentially two different classes of experimental results: the single particle momentum spectra ($dn^-/dy, T_{\text{eff}}$), and the two pion Bose-Einstein correlations ($R_s, R_o, R_l, R_{ol}, \lambda_{\text{dir}}$). We have calculated $\langle f \rangle(p_T, y)$ for the S-S, S-Cu, S-Ag, S-Au, S-Pb and Pb-Pb data from the experiments NA35 [2], NA49 [3], NA44 [4], and for the π -p data from the NA22 experiment [5] at CERN-SPS. From the results for $\langle f \rangle$ as a function of p_T , presented in Fig. 1, one may conclude:

1. universal phase-space density

All the nuclear collision data from the SPS in Fig. 1 are indistinguishably similar, in spite of a factor of ~ 10 difference in multiplicity density.

2. agreement with Bose-Einstein distribution

Using simultaneously pion spectra and pion correlations NA49 has disentangled thermal motion from collective expansion [3]. Taking the NA49 result for the local freeze-out temperature $T = 120$ MeV as the **only** parameter in Eq. (1) one indeed finds good agreement with the data. This is consistent with the thermal nature of the pion source, and inconsistent with the presence of a hypothetical pion condensate.

3. radial flow

Looking in more detail, one finds that the data indicate a somewhat slower decrease with increasing p_T than the Bose-Einstein curve. This is most likely due to radial collective expansion which adds extra transverse momentum to particles, i.e. the local $\langle f \rangle$ values appear in the measurement at a p_T that is higher than the local $\langle p_T \rangle$ in the source reference frame. A detailed study is under way.

4. rapidity dependence

A certain departure from the universal scaling is seen for the data at rapidities close to the projectile rapidity, both at AGS and SPS; moreover, the two results are consistent.




Figure 1: Phase-space density as a function of p_T for different data sets. Heavy-ion data from SPS are indistinguishably similar, although they span over an order of magnitude range in multiplicity. A Bose-Einstein function (Eq. (1)) is superimposed with the three choices of the local freeze-out temperature: 80 MeV, 120 MeV and 180 MeV.

5. high temperature decoupling in $\pi - p$ collisions

In contrast to freeze-out in nuclear collisions which takes place in two steps (chemical at $T \simeq 170$ -180 MeV and thermal at $T \simeq 120$ MeV), pion production in π -p collisions [5] is essentially immediate, without the second evolution stage, and therefore freeze-out temperatures of around 180 MeV should be expected. The data [5] are indeed consistent with this expectation, as seen in Fig. 1.

References

- [1] G.F. Bertsch, Phys. Rev. Lett. **72** (1994) 2349, and **77** (1996) 789e.
- [2] NA35, T. Alber et al., Z. Phys. **C66** (1995) 77, Phys. Rev. Lett. **74** (1995) 1303.
- [3] NA49, H. Appelshäuser, Ph.D. Thesis, Univ. Frankfurt, 1997. H. Appelshäuser et al., Eur. Phys. J. **C2** (1998) 661.
- [4] NA44, H. Beker et al., Phys. Rev. Lett. **74** (1995) 3340. I.G. Bearden et al., Phys. Rev. **C58** (1998) 1656.
- [5] N.M. Agababyan et al., Z. Phys. **C71** (1996) 405.
- [6] E877, J. Barrette et al., Phys. Rev. Lett. **78** (1997) 2961.